

Knowledge and Practices of Physicians about Radiation Protection of Patients during Prescription of Computed Tomography (CT)scan Procedures in Saudi Arabia 2024

Faisal Ahmed Ali Almalki¹, Lujain Jamil Ghafouri², Abulaziz Saud Abdullah alqahtani³, Yasser Mohammed Halawi⁴, Jawhara Abdulaziz Halabi⁵, Ebtisam Rashid Aleshly⁶, Hamad Abdullah Saud Alkathlan⁷, Aeshah Ibrahim Essa Alkhaibari⁸, Fahad Jazi Albogame⁹, Malek Saqr Almutairi¹⁰, Mazen Mushhan Saad Almutairi¹¹

¹CT Scan Specialist, King Abdulaziz Hospital in Jeddah, Saudi Arabia.

²Radiographer, Maternity and children hospital, Saudi Arabia.

³Radiology Technician, Rawda Hospital, Al-Ardh, Saudi Arabia.

⁴Radiology technician, Jeddah Directorate of Health Affairs, Saudi Arabia.

⁵Radiology technician, Forensic Medical Services Center, Saudi Arabia.

⁶X ray technician, Maternity & Children's specialty Hospital, Saudi Arabia.

⁷Radiography Technician, Samtah General Hospital, Saudi Arabia.

⁸Radiological Technology, Maternity and children hospital, Saudi Arabia.

⁹Radiology Technician, Al Masif Health Center, Saudi Arabia.

¹⁰Radiology Technician, Primary Health Care Center, Al Wadi District, Riyadh, Saudi Arabia.

¹¹Radiology Technician, Haremla Health Center, Saudi Arabia.

Received: 13.08.2024

Revised: 19.09.2024

Accepted: 24.10.2024

ABSTRACT

Background: Radiation safety is a concern for patients, physicians, and staff in many departments, including radiology, interventional cardiology, and surgery. Radiation emitted during fluoroscopic procedures is responsible for the greatest radiation dose for medical staff. Radiation from diagnostic imaging modalities, such as computed tomography, mammography, and nuclear imaging, are minor contributors to the cumulative dose exposures of healthcare personnel. However, any radiation exposure poses a potential risk to both patients and healthcare workers alike.

The aims of study: To evaluate the knowledge and practices of physicians in terms of radioprotection of patients when prescribing computed tomography (CT) procedures.

Methods: A cross-sectional study in which a questionnaire with 23 multiple-choice questions was sent to prescribers of CT examinations such as radiotherapists (RMs), other medical specialists (OMSs), general practitioners (GPs) and residents/interns (R/Is). The first eight questions asked about the demographics of the participants, while the remaining questions asked about knowledge of ionizing radiation examinations, doses received, relative risks, and patient radiation safety training.

Results: 223 physicians in all took part in a survey. The reference group, radiation therapists, knew more about irradiating and non-irradiating imaging than the other groups ($P \leq 0.003$). In contrast to the other groups, 67% of the reference group stated that they consider the number of scans the patient has had in the past year ($P \leq 0.002$). Additionally, just 2% of the various groups had a worldwide understanding of the dangers of ionizing radiation exposure ($P \leq 0.73$). Just 12% of the participants, regardless of their seniority or area of expertise, told the patient about the hazards of X-rays at the time of prescription. Finally, only 21% of the participants declared having had training in radiation protection, with no significant differences between the subgroups ($P \leq 0.832$).

Conclusions: The results are comparable to those of earlier research. They demonstrate that prescribers are not well-informed on the dangers of CT exams. Physicians' ongoing professional development should be strengthened, and interns' introductory curriculum should include training on patient radiation protection.

Keywords: CT, Ionizing radiation, Radiation protection, X-rays, Radiation risks

INTRODUCTION

In order to decrease the negative consequences of ionizing radiation, radiation shielding attempts to avoid needless radiation exposure⁽¹⁾. Ionizing radiation is now a necessary instrument in the medical sector for the diagnosis and treatment of a wide range of illnesses. The total lifetime radiation doses that patients and

healthcare professionals get have changed along with its use⁽²⁾. Fluoroscopic imaging, which employs X-rays to produce dynamic and cinematic functional imaging, is the primary source of radiation exposure in medical settings. Patients and medical personnel are exposed to less radiation when they receive formal radiation protection training⁽³⁾.

However, applying radiation safety guidelines can be difficult process, and many interventionalist do not receive formal training in either residency or fellowship on radiation dose reduction⁽⁴⁾. In particular, there is a lack of compliance with radiation safety regulations, especially among physicians or medical personnel that employ fluoroscopic imaging outside of specialized radiology or interventional departments⁽⁵⁾. Numerous fields, such as vascular surgery, gastroenterology, orthopedics, urology, interventional radiology, and interventional cardiology, use fluoroscopy. A comprehensive understanding of radiation exposure dangers and dose reduction strategies will be crucial as radiation exposure increases in frequency⁽³⁻⁵⁾.

Additionally, a significant role in the patient's life is played by medical imaging. It is essential for both diagnosing illnesses and tracking how treatment outcomes change over time. Ionizing radiation (IR) based medical imaging is extensively utilized⁽⁶⁾. The third radiation protection principle limitation does not apply to patients in the medical field⁽⁷⁾. However, more rigorous application of the first two principles justification and optimization is required⁽⁸⁾. The International Commission on Radiological Protection (ICRP) first proposed the idea of diagnostic reference level (DRL) in 1996 to help imaging specialists apply the principle of optimization for the use of ionizing radiation on their patients⁽⁶⁻⁸⁾.

The risk associated with the use of imaging for diagnostic purposes would potentially increase the radiation dose received by patients. Indeed, exposure to IR for diagnostic purposes accounts for 90% of the total dose received by the UK population⁽⁹⁾. Furthermore, in the USA, public exposure to IR has increased sevenfold due to medical imaging during the years 1980–2006. There has also been a more than 100% increase in the number of patients receiving both very high annual doses from IR (>50 mSv) and low doses (<20 mSv)⁽¹⁰⁾. Studies have estimated that the uncertainties associated with low dose levels induced by medical imaging are large^(11, 12).

These results seem to be contradictory with other more recent studies. Indeed, Preston et al., (2007)⁽¹³⁾ in a study on the occurrence of solid cancers in a population exposed to radiation from the Hiroshima atomic bomb, noted that there was a linear increase in the relative risk of cancer for doses between 0 and 2 Gy. They also reported some flattening of the relative cancer risk at higher doses⁽¹³⁾.

In addition, they found a statistically significant dose response when the analyses were restricted to cohort members who received doses of 0.15 Gy or less⁽¹³⁾. Its risks also showed significant variations by gender, attained age, and age at exposure^(13, 14). In addition, other studies have estimated that low-dose IR exposure during medical imaging could cause harm and account for up to 2% of cancers in the US in the future⁽¹⁵⁻¹⁷⁾. In the UK, 100 to 250 cases of death occur each year due to radiological exposures^(18, 19).

Recently, concerns about physicians' awareness of the ionizing radiation exposure dose during diagnostic radiological procedures are increasing^(20, 21). Therefore, it is essential that physicians pay particular attention to the dose delivered to the patient when prescribing the radiological imaging examination. One study has shown that radiation dose awareness among radiologists is insufficient and among non-radiologists is dramatically low⁽²²⁾. In general, various assessments indicate that physicians have low to moderate levels of knowledge about ionizing radiation exposure doses and the expected health risks to the patient⁽²³⁻²⁷⁾.

In Saudi Arabia, 51 nuclear medicine centers operate under the Ministry of Health, governmental sectors, and private hospitals⁽²⁸⁾. These centers conduct approximately 37,655 general nuclear medicine investigations and 12,387 cardiac scans annually. According to a 2018 survey, the country is equipped with 21 positron emission tomography/computed tomography (PET/CT) machines, 55 single-photon emission computed tomography/computed tomography (SPECT/CT) machines, and 35 SPECT and gamma cameras⁽²⁸⁾. By minimizing unnecessary radiation exposure, radiation protection seeks to lessen the negative effects of ionizing radiation. Therefore, this study aimed to evaluate the knowledge and practices of physicians in the field of patient radiation protection during the prescription of CT procedures.

METHODS

This is a retrospective descriptive-analytical study carried from January to May 2024, involving radiotherapists (RMs), other medical specialists (OMSs), general practitioners (GPs) and residents/ interns (R/Is) practicing in Saudi Arabia. In order to evaluate the physician's knowledge related to X-ray exposure and radiation protection practices in CT, an anonymous standardized questionnaire with 3 sections and a total of 23 questions was developed with reference to the literature^(29, 30). It was created on the platform (Google form) and then sent to physicians via a social network linked to their email address. This questionnaire contains 3 sections and a total of 23 questions.

The first section focused on the socio-professional and personal characteristics of the participants: age, gender, practice area, status, and length of professional experience. The second section consisted of 16 questions concerning knowledge of radiation protection: characteristics of X-rays, frequency of use of CT examinations, the use of the Medical Imaging Examination Guide prior to prescription, information to the patient on the

possible risks of such exposure, knowledge of the relative risks of exposure to ionizing radiation during CT procedures, and the tissue most sensitive to damage by ionizing radiation. The third section was devoted to whether or not they had received basic radiation safety training, patient radiation safety training, and whether they felt the need for such training.

Radiotherapists were considered as the reference group for the comparative study with the other groups. Descriptive statistics were produced for the characteristics of the participants, namely age and sex, seniority, sector of activity and status. The categorical variables are expressed as percentages and the participants' scores were classified according to their level of professional experience. To compare responses between the four groups of prescribers, Fisher exact test of the statistical package for the social sciences (SPSS version 28.0) were used. The value $P < 0.05$ indicated the difference was statistically significant.

RESULTS

Table (1) showed Socio-professional characteristics of the study population. In total, two hundred and twenty-three questionnaires were collected and analyzed. General practitioners represented more than a quarter of the participants with 28.2%, followed by interns and residents with 26.5%, then other specialists with a percentage of 23.3%, and finally radiotherapists who represented 22.0%. Furthermore, 28.85% of the specialists exercise in the private sector, 51.92% in the public sector and 19.23% in university hospital centers. There was a slight female predominance (38.1% men versus 61.9% women with a sex ratio of 0.615). Sixty-five participants were over 45 years of age (29.1%), whereas participants under 25 years of age represented only 6.3% and were in their majority interns. Indeed, about 52.0% of the participants had less than 10 years of professional experience, mainly among radiotherapists, residents and interns, whereas 32.7% of the specialists and 44.4% of the generalists had more than 20 years of experience.

Table 1: Demographic data of physicians consulted in this study

Items	MRs	OMSs	GPs	R/Is	Total	P-value
Total	49	52	63	59	223	
Age range (years old)						<0.001
< 35	38(77.6)	3 (5.8)	10(15.9)	50(84.7)	101(45.3)	
≥ 35	11(22.4)	49(94.2)	53(84.1)	9(15.3)	122(54.7)	
Sex						0.210
Male	16(32.7)	25(48.1)	26(41.3)	18(30.5)	85(38.1)	
Female	33(67.3)	27(51.9)	37(58.7)	41(69.5)	138(61.9)	
Sector of activity						<0.001
Private	1 (2.0)	15(28.8)	28(44.4)	1 (1.7)	45(20.2)	
Public	8(16.3)	27(51.9)	33(52.4)	5 (8.5)	73(32.7)	
University Hospital Center	40(81.6)	10(19.2)	2 (3.2)	53(89.8)	105(47.1)	
Seniority (years old)						<0.001
< 10	43(87.8)	10(19.2)	15(23.8)	48(81.4)	116(52.0)	
≥ 10	6(12.2)	42(80.8)	48(76.2)	11(18.6)	107(28.0)	

Note: MRs: radiotherapists; OMSs. Other medical specialists; GPs: General practitioners; R/Is: residents or interns.

Table (2) shows the proportions of correct answers for the questions by group asked for the clinicians in this study. Concerning the distinction between irradiating and non-irradiating examinations, the correct response rate of our prescribers ranged from 41.3% for coronary angiography to 100% for ultrasonography without significant difference between the four groups. For the irradiating examinations, it was 41.3%, 53.8%, 70.0%, 77.1% and 82.9%, respectively for coronary scintigraphy, PET scan, radiography and CT scan. For non-irradiating examinations, the correct response rate of our clinicians was 72.2% for MRI and 100% for ultrasound.

The majority of our participants were prescribers of CT scans, 95.1%, with no significant differences between the different groups ($P = 0.524$). Regardless of their status, the majority of our clinicians reported that they did not use any good practice guide to establish the indications for imaging examinations and to prescribe the least radiating examination ($P = 0.155$). When prescribing a CT examination, 42.9% of the radio-therapists took into account the benefit/risk ratio related to X-rays, with a non-significant difference with the other groups ($P = 0.426$).

Also less than 12% informed their patients of the risks associated with their exposure to X-rays during CT procedures and the resulting benefit/risk ratio ($P = 0.793$). The number of scans performed by the patient in the last year was only taken into account by 21.5% of our clinicians before ordering the examination, with a significant difference between the group of general practitioners and the other groups ($P = 0.004$). On the other hand, no significant difference ($P = 0.126$) was reported on the patients' interest in the risk of X-ray exposure.

Indeed, 57.8% of our clinicians reported that patients never asked them about X-ray risks with rates between 44.9% and 71.4% depending on the group. When patients asked about the risk of ionizing radiation exposures, 75.4% of prescribing physicians explained the expected risk to them based on the benefit/risk ratio. As a result, 12.1% of the specialists changed their prescriptions compared to 2.0% of the radiotherapists ($P = 0.006$).

Concerning the estimation of dose limits, only 11.2% of our prescribers correctly estimated the dose limit recommended by the International Commission on Radiological Protection (ICRP) for the public (1 mSv), the majority of whom were radiotherapists (44.0%, $P = 0.04$). As for the recommended dose limit for pregnant women (1 mSv), only 6.3% of our practitioners had correctly estimated it, with a slightly higher percentage for radiation therapists (12.2%, $P = 0.190$). The ratio between the dose received by a patient during a thoracic scan and that received during a thoracic radiography was correctly estimated by only 6.3% of our prescribing physicians with a difference in favor of the specialist physicians but which was not significant ($P = 0.054$). For the dose delivered to the patient during an abdominal-pelvic scan, only 11.8% of the prescribers had estimated it correctly (7 and 11 mSv) without significant difference between the four groups.

Concerning the radiosensitivity of organs to ionizing radiation, 70.9% of our prescribers answered that the gonads are the most sensitive human tissue to ionizing radiation without significant difference between the four groups. Only 6.8% of our clinicians systematically requested a serum β -HCG assay when prescribing an abdominal-pelvic scan to a woman of childbearing age. Whereas, 31.1% of our prescribers considered that there were no particular precautions to be taken when prescribing a skull scan to a woman of childbearing age.

For the risk of induction of radiation-induced cancer following a single scan, 64.1% of our prescribers answered that there was no risk, without significant difference between the four groups. On the other hand, only 20.0% of our clinicians confirmed that they had received training in radiation protection without significant difference between the four groups ($P = 0.506$). Finally, 90.1% of our practitioners estimated their need for continuous training sessions in radiation protection ($P = 0.112$).

Table 2: Summary of physicians' answers participating on their radiation protection practices according to the question

Questions	Suggestions	MRs	OMSs	MGs	R/Is	Total	P-value
Q1	Yes	48(98.0)	50(96.2)	60(95.2)	54(91.5)	212(95.1)	0.524
	No	1(2.0)	2(3.8)	3(4.8)	5(8.5)	11(4.9)	
Q2	Yes	8(16.3)	2(3.8)	9(14.3)	10(16.9)	29(13.0)	0.155
	No	41(83.7)	50(96.2)	54(85.7)	49(83.1)	194(87.0)	
Q3	Always	21(42.9)	29(55.8)	39(61.9)	30(50.8)	119(53.4)	0.426
	Sometimes	23(46.9)	17(32.7)	16(25.4)	22(37.3)	78(35.0)	
	Never	5(10.2)	6(11.5)	8(12.7)	7(11.9)	26(11.7)	
Q4	Always	6(12.2)	4(7.7)	1(15.9)	5(8.5)	25(11.2)	0.793
	Sometimes	17(34.7)	17(32.7)	25(39.7)	26(44.1)	85(38.1)	
	Never	11(22.4)	15(28.8)	13(20.6)	11(18.6)	50(22.4)	
	If requested	15(30.6)	16(30.8)	15(23.8)	17(28.8)	63(28.3)	
Q5	Always	11(22.4)	13(25.0)	22(34.9)	2(3.4)	48(21.5)	0.004
	Sometimes	18(36.7)	16(30.8)	19(30.2)	25(42.4)	78(35.0)	
	Never	20(40.8)	23(44.2)	22(34.9)	32(54.2)	97(43.5)	
Q6	Once a week	4(8.2)	2(3.8)	1(1.6)	1(1.7)	8(3.6)	
	Once a week	6(12.2)	6(11.5)	3(4.8)	7(11.9)	22(9.9)	
	Once a week	17(34.7)	13(25.0)	14(22.2)	20(33.9)	64(28.7)	
	Never	22(44.9)	31(59.6)	45(71.4)	31(52.5)	129(57.8)	
Q7	Change the procedure	1(2.8)	4(12.1)	1(2.6)	0(0.0)	6(4.2)	0.006
	Reassure if no risk	1(2.8)	1(3.0)	0(0.0)	1(2.9)	3(2.1)	
	Talk about the benefit/risk ratio	30(83.3)	20(60.6)	32(82.1)	25(73.5)	107(75.4)	
	Explain that the risk is negligible	4(11.1)	1(3.0)	2(5.1)	0(0.0)	7(4.9)	
	Change the subject	0(0.0)	7(21.2)	3(7.7)	7(20.6)	17(12.0)	
	Ask the Radiologist	0(0.0)	0(0.0)	1(2.6)	7(2.9)	2(1.4)	
Q8	True	11(22.4)	5(9.6)	4(6.3)	5(8.5)	25(11.2)	0.040
	False	38(77.6)	47(90.4)	59(93.7)	54(91.5)	198(88.8)	
Q9	True	6(12.2)	4(7.7)	2(3.2)	2(3.4)	14(6.3)	0.190
	False	43(87.8)	48(92.3)	61(96.8)	57(96.6)	209(93.7)	
Q10	True	4(8.2)	7(13.5)	2(3.2)	1(1.7)	14(6.3)	0.054
	False	45(91.8)	45(86.5)	61(96.8)	58(98.3)	209(93.7)	

Q11	True	8(16.7)	5(9.8)	6(9.5)	7(11.9)	26(11.8)	0.658
	False	40(83.8)	46(90.2)	57(90.5)	52(88.1)	195(88.2)	
Q12	True	38(77.6)	34(65.4)	41(65.1)	45(76.3)	158(70.9)	0.299
	False	11(22.4)	18(34.6)	22(34.9)	14(23.7)	65(29.1)	
Q13	True	4(8.3)	3(5.8)	4(6.3)	4(6.8)	15(6.8)	0.966
	False	44(91.7)	49(94.2)	59(93.7)	55(93.2)	207(93.2)	
Q14	True	20(40.8)	15(28.8)	9(14.3)	25(43.1)	69(31.1)	0.002
	False	29(59.2)	37(71.2)	54(85.7)	33((56.9)	153(68.9)	
Q15	True	20(40.8)	16(30.8)	22(34.9)	22(37.3)	80(35.9)	0.756
	False	29(59.2)	36(69.2)	41(65.1)	37(62.7)	143(64.1)	
Q16							
MRI	True	28(57.1)	41(78.8)	54(85.7)	38(64.4)	161(72.2)	0.003
	False	21(42.9)	11(21.2)	9(14.3)	21(35.6)	62(27.8)	
Radiography	True	39(79.6)	42(80.8)	48(76.2)	43(72.9)	172(77.1)	0.754
	False	10(20.4)	10(19.2)	15(23.8)	16(27.1)	51(22.9)	
Ultrasound	True	49(100.0)	52(100.0)	63(100.0)	59(100.0)	223(100.0)	
	False	0(0.0)	0(0.0)	0(0.0)	0(0.0)	0(0.0)	
CT	True	42(85.6)	41(78.8)	53(84.1)	48(82.8)	184(82.9)	0.815
	False	7(14.3)	11(21.2)	10(15.9)	10(17.2)	38(17.1)	
Coronaryangiography	True	33(67.3)	29(55.8)	31(49.2)	27(45.8)	120(53.8)	0.123
	False	16(32.2)	23(44.2)	32(50.8)	32((54.2)	103(67.3)	
Scintigraphy	True	26(53.1)	25(48.3)	22(34.9)	19(32.2)	92(41.3)	0.077
	False	23(59.2)	27(51.9)	41(65.1)	40(67.8)	131(58.7)	
Mammography	True	26(53.1)	25(48.3)	22(34.9)	19(32.2)	92(41.3)	0.077
	False	23(59.2)	27(51.9)	41(65.1)	40(67.8)	131(58.7)	
PET/CT	True	35(71.4)	36(69.2)	48(76.2)	37(62.7)	156(70.0)	0.440
	False	14(28.6)	16(30.8)	15(23.8)	22(37.3)	67(30.0)	
Q17	Yes	10(20.4)	6(11.5)	11(17.5)	13(22.0)	40(17.9)	0.506
	No	39(79.6)	46(88.5)	52(82.5)	46(78.0)	183(82.2)	

DISCUSSION

Radiological tests have become increasingly important in medical diagnosis over the last 20 years. CT exams are being used more often, and the risks and harm are widely known and contentious. According to the results of this study, only 11% of our prescribing physicians were aware of diagnostic imaging examinations that use ionizing radiation. This result is much lower than that reported in a Canadian study (91%)⁽³¹⁾. In addition, the group of radiation therapists had the highest score and the group of interns and residents the lowest score in terms of knowledge of radio-diagnostic examinations.

The correlation between the knowledge of radiation imaging and the level of education of our prescribers was not significant ($P < 0.07$) in contrast to a similar study that showed a correlation between the medical specialties that used radiation and their level of education⁽³²⁾. It is also surprising that 27.8% of our prescribers thought that magnetic resonance imaging (MRI) is a radiating imaging test (especially in the intern and resident group).

In terms of radiation-induced risk, 57.8% of our prescribers reported that their patients never showed any interest in the risks of X-rays. These results are similar to those reported in the literature. Indeed, in a Canadian study, 50% of the prescribers reported that none of their patients asked about the risk of radiation⁽³¹⁾. On the other hand, less than 12% of our clinicians informed their patients about the X-ray risks to which they would be exposed. This finding is slightly higher than that reported by a Canadian study which has showed that only 8% of patients were informed by their prescribers of the risks of exposure to ionizing radiation during radiological examinations⁽³¹⁾.

Regarding knowledge of the dose limits recommended by the ICRP for the public and pregnant women, less than 20% of our prescribers had correctly recognized them without significant differences between the different groups. In addition, the correct estimation of the ratio between the effective dose received during a thoracic scan and that received during a thoracic X-ray and the average dose delivered during an abdominal-pelvic scan was made respectively by only 11% and 6% by our prescribers. Also, 64% of our practitioners underestimated the potential risk associated with low doses of X-rays, particularly the risk of radiation-induced cancer.

This low awareness of radiation risks is also common for the group of radiotherapists despite the fact that they are traditionally more aware of the effects of ionizing radiation. These results are similar to those reported by several previous similar studies⁽³¹⁻³³⁾. These low levels of knowledge were confirmed by a study of final year medical school students in Norway and also by a recent study of radiographers in Iran, but are lower than those

shown by a recent study of radiographers in the central region of Ghana⁽³⁴⁻³⁶⁾. These low levels of knowledge were also confirmed by a study of final year students at a medical school in Norway. Indeed, 35.55% of the students reported correct answers despite having radiation safety training during their curriculum⁽²⁹⁾. Even more surprisingly, only 20% of our clinicians confirmed that they had undergone patient radiation safety training, including radiation therapists. This result is slightly lower than the 28.65% reported by Saeed et al., (2018)⁽³⁷⁾ in a similar survey of physicians practicing in Saudi Arabia. In this survey, only 60% of the radiation therapists reported having received radiation safety training, yet it is part of their training curriculum from the first year of residency. This training is also integrated in the form of a training module during the first semesters of medical studies in Saudi Arabia.

CONCLUSION

This study shows that the level of knowledge and practices of physicians in terms of radiation protection is insufficient. This should urgently call upon the competent authorities in this field to integrate more in-depth training on radiation protection of patients into the medical education curriculum, as well as to require qualified and accredited training in radiation protection for practicing physicians. Finally, a guide of radiological procedures and others for regular quality control of radiological equipment are needed to optimize diagnostic procedures and prevent adverse events.

REFERENCES

1. Mitchell EL, Furey P. Prevention of radiation injury from medical imaging. *J Vasc Surg.* 2011 Jan;53(1 Suppl):22S-27S.
2. Tsapaki V, Balter S, Cousins C, Holmberg O, Miller DL, Miranda P, Rehani M, Vano E. The International Atomic Energy Agency action plan on radiation protection of patients and staff in interventional procedures: Achieving change in practice. *Phys Med.* 2018 Aug;52:56-64.
3. The 2007 Recommendations of the International Commission on Radiological Protection. ICRP publication 103. *Ann ICRP.* 2007;37(2-4):1-332.
4. Frane N, Megas A, Stapleton E, Ganz M, Bitterman AD. Radiation Exposure in Orthopaedics. *JBJS Rev.* 2020 Jan;8(1):e0060.
5. Hayda RA, Hsu RY, DePasse JM, Gil JA. Radiation Exposure and Health Risks for Orthopaedic Surgeons. *J Am Acad Orthop Surg.* 2018 Apr 15;26(8):268-277.
6. Matityahu A, Duffy RK, Goldhahn S, Joeris A, Richter PH, Gebhard F. The Great Unknown-A systematic literature review about risk associated with intraoperative imaging during orthopaedic surgeries. *Injury.* 2017 Aug;48(8):1727-1734.
7. Zielinski JM, Shilnikova NS, Krewski D. Canadian National Dose Registry of radiation workers: overview of research from 1951 through 2007. *Int J Occup Med Environ Health.* 2008; 21(4):269-75.
8. Hamada N, Fujimichi Y. Classification of radiation effects for dose limitation purposes: history, current situation and future prospects. *J Radiat Res.* 2014 Jul;55(4):629-40.
9. Hart D, Hillier MC, Shrimpton PC. Doses to Patients from Radiographic and Fluoroscopic X-Ray Imaging Procedures in the UK. HPA-CRCE034. Chilton: Health Protection Agency; 2012.
10. National Council on Radiation Protection and Measurements. Ionizing radiation exposure of the general population of the United States. NCRP Report 160. <http://ncrponline.org/publications/reports/ncrp-report-160/>.
11. Health Physics Society, Position Statement of the Health Physics Society. Radiation risk in perspective. PS010-2. http://hps.org/documents/risk_ps010-2.pdf.
12. American Association of Physicists in Medicine. AAPM Position Statement on Radiation Risks from Medical Imaging Procedures. <https://www.aapm.org/org/policies/details.asp?id%4318&type%4PP¤t%4true>
13. Preston DL, Ron E, Tokuoka S, et al. Solid cancer incidence in atomic bomb survivors: 1958-1998. *Radiat Res.* 2007;168(1):1-64. <https://doi.org/10.1667/RR0763.1>.
14. Committee to Assess Health Risks from Exposure to Low Levels of Ionizing Radiation. Health Risks from Exposure to Low Levels of Ionizing Radiation: BEIR VII Phase 2. Washington, District of Columbia: National Academies Press; 2005.
15. Brenner DJ, Hall EJ. Computed tomography—an increasing source of radiation exposure. *N Engl J Med.* 2007;357(22):2277-2284. <https://doi.org/10.1056/nejmra072149>.
16. Berrington de González A, Mahesh M, Kim KP, et al. Projected cancer risks from computed tomographic scans performed in the United States in 2007. *Arch Intern Med.* 2009;169(22):2071-2077. 10.1001.
17. Hobbs JB, Goldstein N, Lind KE, et al. Physician knowledge of radiation exposure and risk in medical imaging. *J Am Coll Radiol.* 2018;15(1 Pt A):34-43. <https://doi.org/10.1016/j.jacr.2017.08.034>.
18. Einstein AJ, Henzlova MJ, Rajagopalan S. Estimated risk of cancer with radiation exposure from 64-slice computed tomography coronary angiography. *JAMA.* 2007; 298(3):317-323.

- <https://doi.org/10.1001/jama.298.3.317>.
19. Shiralkar S, Rennie A, Snow M, et al. Doctors' knowledge of radiation exposure: questionnaire study. *BMJ*. 2003;327(7411):371–372. <https://doi.org/10.1136/bmj.327.7411.371>.
 20. Naderi M, Salehi F, Maleki S, et al. The need to increase patient awareness of radiation exposure in imaging modalities: a study on the awareness and attitude of patients. *J Med ImagRadiat Sci*. 2021;52(3):450–455. <https://doi.org/10.1016/j.jmir.2021.05.006>.
 21. Almohiy H. Knowledge and awareness of ionizing radiation risks among Saudi obstetricians. *J Radiat Res Appl Sci*. 2020;13(1):542–545. <https://doi.org/10.1080/16878507.2020.1762528>.
 22. Lee RK, Chu WC, Graham CA, et al. Knowledge of radiation exposure in common radiological investigations: a comparison between radiologists and non-radiologists. *Emerg Med J*. 2012;29(4):306–308. <https://doi.org/10.1136/emermed-2011-200481>.
 23. Heyer CM, Hansmann J, Peters SA, et al. Paediatrician awareness of radiation dose and inherent risks in chest imaging studies—a questionnaire study. *Eur J Radiol*. 2010; 76(2):288–293. <https://doi.org/10.1016/j.ejrad.2009.06.014>.
 24. Thomas KE, Parnell-Parmley JE, Haidar S, et al. Assessment of radiation dose awareness among pediatricians. *PediatrRadiol*. 2006;36(8):823–832. <https://doi.org/10.1007/s00247-006-0170-x>.
 25. Arslanoglu A, Bilgin S, Kubal Z, et al. Doctors' and intern doctors' knowledge about patients' ionizing radiation exposure doses during common radiological examinations. *Diagn Interv Radiol*. 2007;13(2):53–55.
 26. Shiralkar S, Rennie A, Snow M, et al. Doctors' knowledge of radiation exposure: questionnaire study. *BMJ*. 2003;327(1):371–372. <https://doi.org/10.1136/bmj.327.7411.371>.
 27. Lee CI, Haims AH, Monico EP, et al. Diagnostic CT scans: assessment of patient, physician, and radiologist awareness of radiation dose and possible risks. *Radiology*. 2004;231(2):393–398. <https://doi.org/10.1148/radiol.2312030767>.
 28. Current situation of nuclear medicine in Saudi Arabia. Alqarni A, Farghaly H, Nasr H. *Int J Sci Res*. 2019;8
 29. Kada S. Awareness and knowledge of radiation dose and associated risks among final year medical students in Norway. *Insights Imaging*. 2017;8(6):599–605. <https://doi.org/10.1007/s13244-017-0569-y>.
 30. Yurt A, Çavuşoğlu B, Günay T. Evaluation of awareness on radiation protection and knowledge about radiological examinations in healthcare professionals who use ionized radiation at work. *Mol Imaging Radionucl Ther*. 2014;23(2):48–53. <https://doi.org/10.4274/mirt.00719>.
 31. Ricketts ML, Baerlocher MO, Asch MR, et al. Perception of radiation exposure and risk among patients, medical students, and referring physicians at a tertiary care community hospital. *Can Assoc Radiol J*. 2013;64(3):208–212. <https://doi.org/10.1016/j.carj.2012.05.002>.
 32. O'Sullivan J, O'Connor OJ, O'Reagan K, et al. An assessment of medical students' awareness of radiation exposures associated with diagnostic imaging investigations. *Insights Imaging*. 2010;1(2):86–92. <https://doi.org/10.1007/s13244-010-0009-8>.
 33. Gervaise A, Esperabe-Vignau F, Pernin M, et al. Evaluation of the knowledge of physicians prescribing CT examinations on the radiation protection of patients. *J Radiol*. 2011;92(7-8):681–687. <https://doi.org/10.1016/j.jradio.2011.03.023>.
 34. Kazemi Z, Hajimiri K, Saghatchi F, et al. Assessment of the knowledge level of radiographers and CT technologists regarding computed tomography parameters in Iran. *Radiat Med Protect*. 2023;4(1). <https://doi.org/10.1016/j.radmp.2023.01.002>. in press.
 35. Fiagbedzi E, Gorleku P, Nyarko S, et al. Assessment of radiation protection knowledge and practices among radiographers in the central region of Ghana. *Radiat Med Prot*. 2022;3(3):146–151. <https://doi.org/10.1016/j.radmp.2022.06.001>.
 36. Riz HE, Frush DP, Harker MJ, et al. Peer assessment of pediatric surgeons for potential risks of radiation exposure from computed tomography scans. *J Pediatr Surg*. 2007;42(7):1157–1164. <https://doi.org/10.1016/j.jpedsurg.2007.02.007>.
 37. Saeed MK, Al-shaari H, Almarzooq MMS, et al. Radiation awareness among physicians about the dangers of radiological examinations on the health of workers and their patients in Saudi Arabia. *J Radiat Res Appl Sci*. 2018;11(4):299–304.